Revisiting Electron Pathways in Photosynthesis: A Proposal for a Delayed-Choice Quantum Eraser Experiment in Bacterial Reaction Centers

Douglas C. Youvan

doug@youvan.com

October 27, 2024

The photosynthetic reaction center (RC) in bacteria serves as a remarkably efficient system for converting light into chemical energy through highly controlled electron transfer pathways. Despite its structural symmetry, electron transfer in the bacterial RC occurs almost exclusively along the A-branch, while the B-branch remains largely inactive. This pathway selectivity has been traditionally attributed to structural and energetic differences between the branches. However, recent discoveries in quantum biology suggest that quantum coherence might play a role in such biological processes, raising questions about whether measurement itself influences pathway determination in the RC. This paper proposes a delayed-choice quantum eraser experiment to explore this possibility. By introducing a quantum eraser pulse after electron transfer begins, we aim to disrupt which-path information, potentially revealing a superposition state across both branches. Observing interference patterns would imply that quantum coherence affects pathway selectivity, challenging conventional views of biological determinism in photosynthetic processes and opening new pathways in quantum biology.

Keywords: photosynthesis, reaction center, electron pathways, quantum coherence, quantum eraser, delayed-choice experiment, bacterial photosynthesis, quantum biology, interference, pathway selectivity.

Abstract

This paper introduces a novel experimental approach to investigate the underlying mechanism of electron transfer asymmetry in the bacterial photosynthetic reaction center (RC) through a delayed-choice quantum eraser design. In typical bacterial RCs, electron transfer occurs predominantly along the A-branch, with the B-branch remaining largely inactive. This asymmetry has been widely attributed to structural factors, such as protein environments and molecular energy gradients, which create a natural bias favoring the A-branch. However, the possibility that measurement-induced collapse could influence pathway selectivity has not been explored in depth. In this study, we propose an experiment to determine if measurement plays a role in pathway asymmetry by introducing a quantum eraser pulse that disrupts which-path information after the electron has initiated transfer along the A-branch. By "erasing" this which-path information, we seek to induce a superposition state in which the electron could be observed in both the A- and B-branches, revealing interference patterns if coherence is preserved between the pathways.

The implications of this experiment reach beyond photosynthetic electron transfer, as observing interference would suggest that biological systems might exploit quantum coherence on ultrafast timescales, retaining multiple pathway possibilities until measured. This phenomenon would challenge the current understanding of intrinsic pathway selectivity in photosynthetic RCs and could lead to a broader paradigm for understanding quantum mechanics within biological processes. Should interference be detected, it would imply that pathway determination may not be solely intrinsic but could be influenced by the act of observation itself, much like systems in quantum mechanics where outcomes remain undecided until measurement. Such results could reveal new dimensions of quantum behavior within biochemical systems, potentially influencing fields like quantum biology, biophysics, and even quantum information science. The experimental design, methodology, and anticipated challenges, including coherence maintenance and detection sensitivity, are discussed to provide a comprehensive framework for exploring these foundational questions.

1. Introduction

Background on Photosynthetic Reaction Centers

Photosynthetic reaction centers (RCs) are pivotal in the energy conversion processes of photosynthetic organisms, where they serve as the sites of initial light energy capture and conversion into chemical energy. In bacterial RCs, this energy conversion process begins when absorbed photons excite specialized pigments, triggering an electron transfer sequence that ultimately generates a proton gradient across the membrane. This gradient drives ATP synthesis, providing the essential energy for cellular activities. The electron transfer sequence in these bacterial RCs is highly efficient, ensuring minimal energy loss through a finely controlled pathway.

The bacterial RC is asymmetric in its structure, containing two nearly identical electron transfer pathways known as the A- and B-branches. Despite this structural symmetry, electron transfer occurs predominantly through the A-branch, with the B-branch remaining largely inactive under standard conditions. This selectivity has been attributed to small structural differences and subtle variations in protein environments, which collectively create an energetic landscape favoring the A-branch. For instance, differences in distance, orientation, and local electric fields within the protein complex contribute to a lower energy barrier for electron transfer along the A-branch. While structural explanations for pathway asymmetry are widely accepted, it remains unclear whether this preference is strictly a result of physical configuration or if it may also be influenced by quantum mechanical effects.

Quantum Mechanical Influence in Photosynthesis

The role of quantum mechanics in biological systems has gained increasing attention, particularly in photosynthesis, where quantum coherence may play a role in optimizing energy transfer. Recent studies have demonstrated that exciton dynamics—interactions between photons and electron-hole pairs—within photosynthetic systems can exhibit coherence effects on ultrafast timescales, suggesting that quantum effects might help enhance the efficiency of light capture and energy transfer. Observations of coherence suggest that components of the photosynthetic machinery can maintain quantum superpositions long

enough to influence the pathway of energy transfer, thus challenging the classical notion of discrete, deterministic pathways.

In the context of the bacterial RC, the observation of coherence has led to speculation that the electron transfer asymmetry might not be entirely intrinsic to the protein's structure. Instead, it might be partially influenced by the collapse of a quantum state at the moment of measurement, implying that pathway selection could be affected by the act of observing the electron's position. This parallels delayed-choice experiments in quantum mechanics, where particles appear to “choose” their path only upon measurement. If electron pathway selection in bacterial RCs operates similarly, it would mean that the system might retain a superposition state over both A- and B-branches until a collapse is induced by observation.

Proposing a Quantum Eraser Approach

In light of these considerations, we propose an experiment using a delayed-choice quantum eraser approach to investigate the pathway asymmetry in bacterial RCs. The delayed-choice quantum eraser experiment would involve first initiating electron transfer along the A-branch and then introducing an eraser pulse to potentially “erase” which-path information. This pulse would be timed to coincide with electron localization within an intermediary molecule, allowing us to examine if pathway determination can be reversed or influenced after the transfer process has begun.

Our objective is to assess whether the electron transfer dynamics in the bacterial RC can reveal superposition states involving both A- and B-branches. The presence of an interference pattern would suggest that the system retains quantum coherence in its electron transfer pathways, implying that the observed asymmetry may arise not solely from structural constraints but potentially as an artifact of measurement. If successful, this experiment would challenge our current understanding of biological pathway selectivity, providing evidence that quantum effects in photosynthetic systems might allow biological structures to maintain coherence until measurement collapses the system into a specific pathway.

By extending concepts from quantum mechanics to biological electron transfer processes, this study aims to open new avenues in the field of quantum biology.

Our findings could have broad implications for how quantum mechanics may underpin biological functions and how measurement may play an active role in defining biological outcomes in systems where coherence could enhance efficiency and adaptability. Through this delayed-choice quantum eraser experiment, we hope to provide insights that not only deepen our understanding of photosynthetic electron transfer but also challenge the boundaries between classical and quantum descriptions of biological systems.

2. Experimental Design and Methodology

This section details the experimental setup for applying a delayed-choice quantum eraser approach to electron transfer pathways in bacterial photosynthetic reaction centers (RCs). The design seeks to determine whether electron transfer asymmetry in the RC is due to structural constraints or if quantum coherence effects allow both A- and B-branches to maintain superposition states until observation.

2.1. Reaction Center Sample Preparation

To preserve the natural electron transfer dynamics of the bacterial photosynthetic reaction center, precise sample preparation is crucial:

- **Isolation of Bacterial Reaction Centers**: Bacterial RCs, particularly from *Rhodobacter sphaeroides*, are ideal due to their well-characterized structure and consistent electron transfer pathways. The RCs are isolated using biochemical techniques, such as detergent solubilization and chromatographic purification, ensuring minimal structural alteration.
- **Embedding Medium**: To maintain both branches' function and facilitate electron transfer coherence, the RCs are embedded in a stabilizing medium. This medium should ideally mimic the native environment of the RC while reducing environmental noise that could cause premature decoherence. Techniques such as encapsulating RCs in lipid bilayers or specific polymer matrices can help sustain the delicate structural and functional integrity needed for quantum coherence over the experiment’s duration.

- **Cryogenic or Room-Temperature Conditions**: Given that quantum coherence can be more readily maintained at lower temperatures, the experiment could benefit from being conducted at cryogenic temperatures, balancing the need for coherence preservation with biologically relevant conditions.

2.2. Pump-Probe Excitation for Pathway Initiation

The pump-probe technique allows the precise initiation and tracking of electron transfer events, capturing real-time dynamics and initial pathway selection:

- **Initial Pump Pulse**: A femtosecond laser pulse is used to excite the RC's primary electron donor (typically the special pair of bacteriochlorophyll molecules). This excitation triggers electron transfer down the A-branch under typical conditions, establishing which-path information early in the transfer process.
- **Time-Resolved Spectroscopy for Real-Time Probing**: To observe the electron's initial localization, time-resolved pump-probe spectroscopy is employed. After the pump pulse, a series of probe pulses is applied to measure transient absorption or fluorescence changes, providing a time-resolved map of electron movement within the RC. This setup enables us to confirm that electron transfer predominantly follows the A-branch, establishing an initial baseline for which-path information.
- **Verification of Pathway Directionality**: Real-time probing provides critical confirmation of pathway directionality by capturing electron presence within specific molecular intermediates in the A-branch. This step establishes the electron's preference for the A-branch before the introduction of the eraser pulse.

2.3. Introduction of the Quantum Eraser Pulse

The quantum eraser pulse is a secondary laser pulse introduced to interfere with the initial which-path information, potentially resetting the pathway determination and allowing a superposition state across both branches.

- **Timing and Nature of the Delayed Eraser Pulse**: The eraser pulse is applied shortly after detecting initial electron transfer down the A-branch, timed to coincide with the electron's presence within intermediary molecules common to both branches. This pulse must be carefully tuned in both timing and energy to maximize coherence across the A- and B-branches without disrupting the RC's structure.
- **Purpose of the Eraser Pulse**: The aim is to induce a coherent state that "erases" the initial which-path information, allowing the electron to explore both branches simultaneously in a superposition. In effect, this pulse should interfere with any classical localization of the electron on the A-branch, potentially merging it with the B-branch, thereby reestablishing a state of quantum uncertainty.
- **Eraser Pulse Wavelength and Power**: The wavelength and power of the eraser pulse are selected based on the specific absorption characteristics of the RC intermediates to ensure efficient coupling. The energy should be sufficient to induce coherence without exciting the RC beyond the levels necessary for electron transfer.

2.4. Detection of Interference Patterns

The final step is to monitor for interference patterns, which would indicate whether the electron transfer maintains quantum coherence between the A- and B-branches.

- **Fluorescence or Oscillatory Detection in Acceptor Molecules**: Upon the completion of electron transfer, fluorescence or transient absorption measurements are used to track electron localization in the acceptor molecules of both branches. The presence of oscillatory or interference patterns would reveal coherence effects, suggesting that the electron was in a superposition across the two branches.
- **Interference as a Signal of Quantum Coherence**: Observing interference patterns would suggest that both pathways can retain coherence, implying that the A-branch preference might result from the measurement collapse rather than an intrinsic property. This interference could manifest as

oscillatory changes in fluorescence intensity or specific spectral features characteristic of coherent states.

- **Control Experiments**: To ensure results are due to the quantum eraser pulse and not structural effects, control experiments would involve repeating the pump-probe sequence without the eraser pulse. Lack of interference patterns in the control would confirm that observed interference in the main experiment results from the introduced eraser pulse and not inherent symmetry breaking in the RC.
- **Signal Interpretation**: The presence of interference would indicate that pathway asymmetry is not solely due to protein structure but might result from measurement effects, as seen in quantum systems. An absence of interference, however, would suggest that the RC structure enforces intrinsic pathway selectivity, independent of measurement.

This experimental design leverages cutting-edge techniques in ultrafast laser spectroscopy and quantum coherence control to address fundamental questions about quantum mechanics in biological systems. Through time-resolved detection of electron pathways and the careful application of a quantum eraser pulse, this experiment could provide unprecedented insights into the role of observation in determining biological pathway outcomes, with broad implications for quantum biology and biophysics.

3. Hypothesis and Predictions

This experiment seeks to test whether the observed asymmetry in electron transfer within the bacterial photosynthetic reaction center (RC) is due to intrinsic structural constraints or if it may arise from the act of measurement itself. The hypothesis is grounded in the principles of quantum mechanics, specifically the role of measurement in collapsing superposition states into specific outcomes. In conventional delayed-choice quantum eraser experiments, particles retain superposition properties until observation, at which point they “choose” a path. If this principle applies to the bacterial RC, we hypothesize that the electron may

exist in a superposition across the A- and B-branches until the moment of detection.

Our central hypothesis is that introducing a delayed-choice eraser pulse will interfere with initial which-path information, disrupting the electron's deterministic localization on the A-branch and potentially creating a state of superposition between both pathways. If this hypothesis holds true, then interference patterns should emerge, reflecting a coherent superposition of electron pathways across both the A- and B-branches. Conversely, if pathway selectivity is a result of inherent structural factors within the RC, the coherence between pathways would collapse prior to the eraser pulse's influence, and no interference patterns would be detected.

Predicted Outcomes

1. **Interference Pattern Detection**
    - **Superposition and Quantum Coherence**: If we observe interference patterns in electron localization, this would strongly suggest that the electron initially exists in a superposition state across both the A- and B-branches. This outcome would imply that the observed asymmetry in electron transfer is influenced by measurement, rather than an intrinsic bias. In this case, pathway asymmetry would emerge as an artifact of the experimental setup, where the act of measurement collapses the electron's state into a defined pathway.
    - **Implications for Quantum Coherence in Biology**: Detecting an interference pattern would support the idea that quantum coherence is preserved in biological systems over timescales relevant to biological processes. This would lend credence to theories in quantum biology suggesting that biological structures, like photosynthetic RCs, might exploit quantum superposition to maximize efficiency or adaptability. Such a finding could also imply that other biochemical processes traditionally viewed as classical might involve quantum coherence under certain conditions.

2. **Absence of Interference Patterns**
    - **Intrinsic Asymmetry Due to Structural Factors**: If no interference pattern emerges, it would indicate that electron transfer in the RC is primarily determined by structural and energetic constraints that inherently favor the A-branch. This outcome would suggest that pathway asymmetry is built into the protein structure, with protein environments, distances between cofactors, or energy gradients enforcing a deterministic preference for the A-branch.
    - **Classical Behavior in Electron Pathway Selection**: An absence of interference would support the classical view that biological pathway selection in photosynthetic systems is fixed by structural elements rather than being susceptible to quantum coherence. This result would imply that the electron's pathway is effectively "chosen" by the RC's architecture itself, and the delayed-choice eraser pulse cannot alter this choice due to immediate decoherence.

In either case, the findings from this experiment would have substantial implications. Observing interference would open new pathways for understanding quantum effects in biology, suggesting that quantum mechanics plays a more dynamic role in electron transfer processes than previously thought. On the other hand, the absence of interference would reinforce the view that biological electron transfer pathways are deterministic and rooted in molecular structure, aligning with traditional biophysical interpretations. This experiment, therefore, provides a rigorous test for the potential influence of quantum mechanics in complex biological systems, revealing whether electron pathway selectivity in photosynthetic RCs is a fundamental property or a phenomenon shaped by observation.

4. Significance and Implications

This proposed experiment has profound implications for our understanding of quantum mechanics in biological systems, the nature of biological determinism, and the future direction of research in quantum biology. If successful, the results

could redefine how we interpret pathway selection in complex biological processes, emphasizing the role of quantum phenomena in systems traditionally explained by classical biochemistry.

4.1. Quantum Mechanics in Biological Systems

A successful demonstration of interference effects in the bacterial reaction center (RC) would directly challenge the classical perspective that biological processes are strictly deterministic and governed by structural constraints. If interference patterns are observed, it would suggest that quantum superposition can be maintained in biological electron transfer pathways until actively observed or measured. Such a finding would imply that the RC and possibly other biological systems could preserve coherent quantum states, even in complex environments typically prone to rapid decoherence.

This evidence would reinforce theories suggesting that biological organisms might naturally exploit quantum coherence to achieve functional advantages, such as enhanced efficiency in energy transfer, adaptability, or error correction in biochemical processes. For example, coherence in electron pathways could allow the RC to select the most efficient or energetically favorable route for electron transport, dynamically adjusting based on environmental conditions. Such mechanisms could underlie the remarkable efficiency observed in photosynthesis, where energy transfer approaches nearly perfect efficiency under natural conditions. More broadly, this result would position quantum mechanics as a fundamental component of biological functionality, potentially operating in other biological systems where classical explanations have so far sufficed.

4.2. Revisiting Biological Determinism

The findings from this experiment could provoke a significant re-evaluation of biological determinism in photosynthetic pathways and other biochemical processes. Traditional views hold that structural and energetic factors within the RC strictly determine pathway selectivity, with molecular architecture biasing electron flow down the A-branch in bacterial photosynthetic systems. However, if pathway selection is shown to depend partially on measurement, this would

imply that the RC operates under principles similar to quantum measurement theory, where the pathway is undetermined until an observational collapse.

In this framework, the pathway asymmetry observed in electron transfer might be influenced by the experimental setup itself rather than strictly intrinsic structural properties. Such an insight would challenge a century-old foundation of biophysics, suggesting that pathway selectivity could be dynamic, determined not only by protein structure but also by interactions with external observables, such as light pulses or probe measurements. Consequently, this would shift the paradigm in molecular biology, hinting that observation and measurement might play a more active role in defining biological behavior than previously believed.

Reframing pathway selection as a quantum measurement process could redefine our understanding of "choice" and "decision-making" in biochemical processes, suggesting that some biological phenomena might not be inherently fixed but instead fluctuate based on quantum mechanical states until they are measured or influenced by environmental factors. This approach could open the door to reinterpreting other deterministic biological processes through a quantum lens, with implications that stretch across various fields in biology and chemistry.

4.3. Broader Applications and Future Research

The experimental design and methodology proposed here could inspire a range of similar studies in other biological and biochemical systems where pathway asymmetry and selectivity are observed. For example, enzyme-substrate interactions, ion channel dynamics, or metabolic pathways might also exhibit sensitivity to quantum coherence or measurement effects. If pathway selection in these systems can be influenced by quantum eraser or delayed-choice setups, it would suggest a broad, underlying principle in biology: that some functional pathways might remain in superposition states until influenced by measurement, allowing for probabilistic decision-making within the molecular framework.

Beyond biology, these findings could have significant implications for quantum computing, materials science, and even artificial intelligence. In quantum computing, controlling decoherence is essential, as maintaining superposition states enables parallel computation across multiple states. Observing how

biological systems sustain coherence under noisy conditions could provide valuable insights into designing more resilient quantum devices or methods for error correction. In materials science, understanding how biological systems navigate and exploit quantum coherence could inform the development of novel materials with tunable pathways or dynamic responsiveness based on quantum states.

Moreover, if biological systems inherently exploit quantum effects, it could pave the way for future bio-inspired technologies that leverage quantum principles for optimization and adaptability. For example, synthetic systems designed to mimic photosynthetic RCs could use superposition and coherence to achieve unprecedented efficiency in energy capture and transfer. Additionally, these insights might contribute to the emerging field of quantum biology, where understanding the quantum foundations of life could reveal new aspects of evolution, adaptability, and resilience in living organisms.

Ultimately, the results from this experiment could reshape the way scientists view and study the intersection between quantum mechanics and biology. They might also influence other domains where the quantum-classical boundary is crucial, leading to a holistic framework for investigating how quantum effects influence complex systems, from the cellular level to the macroscopic. This broader perspective could ultimately drive interdisciplinary research aimed at uncovering the role of quantum mechanics in both living and synthetic systems, enriching our fundamental understanding of life and the universe.

5. Experimental Challenges and Feasibility

The experimental design proposed here, though promising, entails significant technical challenges due to the delicate nature of quantum coherence in biological systems and the need for ultrafast precision in measuring electron transfer dynamics. Addressing these challenges will require careful calibration of equipment and methodologies, as well as innovations in detection and preservation techniques to ensure that any observed quantum effects are valid and reproducible.

5.1. Timing and Precision

One of the primary challenges is achieving the femtosecond timing precision necessary to synchronize the eraser pulse with the ultrafast dynamics of electron transfer in the bacterial reaction center (RC). Electron transfer within the RC occurs on a picosecond to femtosecond timescale, and any delayed eraser pulse must be precisely timed to match this rapid transfer process.

- **Laser Synchronization**: The experiment requires two distinct laser pulses—a pump pulse to initiate electron transfer and a precisely timed eraser pulse to reset which-path information. Synchronizing these pulses down to the femtosecond scale demands an advanced laser system capable of minimal jitter and highly accurate timing control. State-of-the-art ultrafast laser systems, potentially with feedback loops to account for timing drift, will be necessary to maintain the precision required for observing quantum interference effects.
- **Delay Control Mechanisms**: Fine-tuning the delay between the pump and eraser pulses is essential. Using optical delay lines with sub-femtosecond resolution can help achieve the exact timing needed to position the eraser pulse during electron transfer in the intermediary molecules. Such precision timing systems, while available, are complex and require regular recalibration to account for environmental fluctuations.
- **Temporal and Spatial Precision in Detection**: The pump and eraser pulses must also be spatially aligned within the reaction center to interact effectively with the electron transfer pathway. Careful alignment and temporal synchronization of detection equipment, such as spectrometers or streak cameras, will be required to capture electron localization events accurately.

5.2. Maintaining Quantum Coherence

Another formidable challenge is the inherent decoherence present in biological systems. Biological environments are generally “noisy” due to thermal fluctuations, molecular interactions, and a high degree of complexity, all of which

can disrupt quantum coherence. To successfully detect interference patterns, the system must remain coherent over the timescales involved in electron transfer.

- **Cryogenic Temperatures**: Maintaining quantum coherence in a biological system at room temperature is exceedingly difficult due to rapid decoherence. Lowering the temperature of the system can significantly reduce environmental noise and help preserve coherence. Cryogenic conditions, which slow down molecular vibrations and other decoherence-inducing processes, could enhance the feasibility of observing quantum effects in the RC. However, balancing cryogenic temperatures with biological integrity is challenging, as temperatures too low could alter the RC's functionality.
- **Specialized Embedding Media**: Another potential method for preserving coherence involves embedding the RCs in media that simulate the natural biological environment while reducing noise. Specialized gels, lipid bilayers, or synthetic matrices could help shield the RC from environmental interactions that lead to decoherence. Selecting or developing media that mimic natural conditions without compromising the electron transfer pathways would be crucial to ensuring accurate results.
- **Isolation Techniques**: Physically isolating the RCs to reduce interaction with external molecules or ions can help mitigate decoherence. Techniques such as embedding RCs within nanoporous materials or enclosing them in isolated microenvironments could also offer additional coherence preservation. The challenge, however, is maintaining functionality in such isolated conditions, as it could potentially interfere with the natural operation of the RC.

5.3. Detecting Weak Interference Signals

Even if quantum coherence is achieved, detecting interference patterns within a biological matrix presents unique difficulties. Interference signals could be subtle and may be masked by the background noise inherent in biological systems, necessitating highly sensitive detection methods.

- **High-Sensitivity Detectors**: Capturing faint interference signals in a biological setting requires detectors with high sensitivity and low noise levels. Advanced photodetectors, such as superconducting nanowire single-photon detectors or avalanche photodiodes, could be employed to improve signal-to-noise ratios. Additionally, using multiple detectors positioned around the RC sample may help capture interference patterns from various angles, providing more comprehensive data on electron localization and pathway superposition.
- **Enhanced Fluorescence Detection**: Since electron transfer events within the RC produce changes in fluorescence, time-resolved fluorescence techniques can be employed to detect interference. Techniques like fluorescence lifetime imaging microscopy (FLIM) or femtosecond fluorescence spectroscopy can offer high temporal resolution, allowing researchers to track fluorescence oscillations indicative of interference patterns. These techniques would require meticulous calibration to separate interference-related signals from background fluorescence.
- **Signal Processing and Noise Reduction**: Advanced signal processing techniques, such as wavelet transformation, machine learning-based filtering, or Fourier analysis, could help extract interference signals from noisy biological data. Employing algorithms designed to detect specific interference oscillations may enhance the clarity of results, making it easier to discern subtle quantum effects within the electron transfer pathway.
- **Control Experiments to Validate Signal Origin**: To confirm that any detected interference pattern results from quantum coherence, control experiments must be conducted. By removing the eraser pulse or altering its timing, researchers can compare outcomes to verify that observed interference is indeed due to the delayed-choice setup and not an artifact of the system. Additionally, conducting the experiment with structurally modified RCs, where the A- or B-branch is selectively deactivated, could help validate the pathway superposition hypothesis.

Overall, the feasibility of this experiment hinges on overcoming these challenges through a combination of cutting-edge technology and precise experimental

control. Advances in ultrafast laser systems, cryogenic techniques, high-sensitivity detection, and signal processing make this experiment technically achievable, albeit with considerable effort and resources. Successfully addressing these challenges would not only validate the proposed experimental design but also open up new avenues for applying quantum mechanics to biological systems, marking a significant step forward in quantum biology.

6. Note Added on Vibronic Coupling

If the 1 picosecond (ps) electron transfer time in the photosynthetic reaction center (RC) is due to vibrational-electronic coupling (also known as vibronic coupling), it suggests that vibrational energy from molecular modes within the RC helps drive the electron transfer by making the process isoenergetic. This means that the electron transfer could proceed more efficiently because it effectively "rides" on a vibrational mode that matches the energy difference between the donor and acceptor states. Here's a closer look at the implications of this mechanism:

How Vibrational-Electronic Coupling Works in Electron Transfer

1. Vibronic Coupling for Energy Matching:
    - In vibronic coupling, electronic states are coupled with specific vibrational modes of the molecular structure, enabling electrons to transfer more easily between states if vibrational energy can bridge any energy gap.
    - If the electron transfer from the special pair to the bacteriopheophytin is isoenergetic, it implies that the system has finely tuned vibrational modes that bring the donor and acceptor into near resonance. This tuning reduces the need for an external energy input, allowing the electron to transfer with high efficiency and speed.

2. Facilitated Transfer on the Picosecond Scale:
    - The picosecond timescale corresponds well with vibrational periods in molecular systems, suggesting that specific vibrations within the RC's molecular structure play a direct role in aligning the energy levels.
    - For instance, vibrational modes on the order of terahertz (THz) frequencies, which correspond to picosecond cycles, could be involved. These modes periodically bring the electron's energy in the donor and acceptor into alignment, facilitating a smooth transfer without requiring large energetic "jumps."
3. Quantum Coherence Enhanced by Vibronic Coupling:
    - Vibronic coupling can help maintain quantum coherence between the pathways (A- and B-branches) by synchronizing the electron transfer with specific vibrational states.
    - If the electron remains coherent with both branches due to resonance conditions created by these vibrational states, it could explain why electron transfer shows such high efficiency along the A-branch. The vibrational states essentially reinforce coherence within the chosen pathway (A-branch), while potentially suppressing coherence in the alternative pathway (B-branch).

Implications for the Eraser Pulse

If vibrational-electronic coupling is indeed at play, the eraser pulse must be designed to interact specifically with both the vibrational modes and electronic states involved in the transfer. Here's how this might affect the specifics of the eraser pulse:

1. Pulse Timing and Phase Matching:
    - The eraser pulse should be timed precisely to interact with the electron transfer when vibrational states are facilitating coherence between pathways. A femtosecond (fs) to picosecond pulse would be necessary to match the vibrational-electronic timescale.

- By tuning the pulse to the vibrational frequency, we could enhance or disrupt the vibronic coherence. This could mean either stabilizing the superposition across pathways or scrambling it, depending on the pulse's phase relationship to the vibrational mode.

2. Frequency and Wavelength Tuning for Vibronic Interactions:
    - The eraser pulse would ideally be tuned to interact with both the electronic and vibrational energy levels specific to the RC's molecular structure. This might involve resonant frequencies close to the electronic absorption of the RC, but with an added consideration for vibrational resonance frequencies that match the vibronic coupling.
    - For instance, if a specific vibrational mode (say, a C=O stretching vibration) is involved in the coupling, the pulse could target the energy region around that vibrational frequency to disrupt or maintain coherence.
3. Selective Interference with Vibronic States:
    - The eraser pulse could act to selectively disrupt vibrational coherence, which would erase the energy alignment between the A- and B-branch pathways. This loss of vibrational coherence would make it harder for the electron to "sample" both pathways coherently, effectively collapsing it into a single, preferred pathway.
    - Alternatively, if designed to preserve coherence, the eraser pulse could be tuned to reinforce the alignment of vibrational modes across both branches, potentially allowing both pathways to contribute to the electron's transfer state in a superposition.
4. Environmental Coupling Considerations:
    - Because vibrational-electronic coupling is sensitive to the environment, the eraser pulse would also need to avoid inducing decoherence by disturbing the RC's delicate surroundings. Therefore, the pulse should be intense enough to influence the vibronic states without causing broader decoherence from environmental interactions.

Potential Experimental Outcomes

- Interference Patterns: If the pulse successfully maintains coherence across both branches, interference patterns in the electron transfer would indicate that the electron had access to both pathways in a superposition, facilitated by the vibrational resonance. This would imply that quantum coherence plays a role in pathway selectivity, possibly driven by vibrational-electronic coupling.
- Loss of Pathway Selectivity: If coherence is disrupted (either by removing vibronic resonance or collapsing the vibrational state), the electron transfer would likely revert to classical behavior, with a clear preference for a single pathway (likely the A-branch, due to traditional structural and energetic biases).

Summary

If the 1 ps transfer time is due to vibrational-electronic coupling, then the eraser pulse should specifically target both the electronic and vibrational states responsible for maintaining coherence across the A- and B-branch pathways. The pulse would need to match the vibrational resonance to either enhance or scramble the quantum coherence enabled by these vibronic interactions. By interacting with the RC at this precise resonance, the experiment could reveal whether quantum coherence and vibronic coupling are indeed responsible for the high efficiency and selectivity observed in the bacterial photosynthetic RC, thus supporting the role of quantum biology in this process.

References

1. **Blankenship, R. E.** (2014). *Molecular Mechanisms of Photosynthesis*. 2nd ed. Wiley-Blackwell.
    - This book provides comprehensive coverage of the molecular processes involved in photosynthesis, including electron transfer in bacterial reaction centers.
2. **Engel, G. S., Calhoun, T. R., Read, E. L., Ahn, T.-K., Mancal, T., Cheng, Y.-C., Blankenship, R. E., & Fleming, G. R.** (2007). Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature*, 446(7137), 782–786.
    - A landmark paper demonstrating quantum coherence in photosynthetic energy transfer, suggesting that quantum effects may contribute to the high efficiency of photosynthetic processes.
3. **Scholes, G. D., Fleming, G. R., Olaya-Castro, A., & van Grondelle, R.** (2011). Lessons from nature about solar light harvesting. *Nature Chemistry*, 3(10), 763–774.
    - Discusses how natural systems use quantum coherence and energy transfer processes, with implications for the efficiency of photosynthetic reaction centers.
4. **Ishizaki, A., & Fleming, G. R.** (2009). Theoretical examination of quantum coherence in a photosynthetic system at physiological temperature. *Proceedings of the National Academy of Sciences*, 106(41), 17255–17260.
    - Provides theoretical analysis of how quantum coherence might be maintained in photosynthetic systems at physiological temperatures, relevant to understanding coherence preservation in biological environments.

5. **Scully, M. O., & Drühl, K.** (1982). Quantum eraser: A proposed photon correlation experiment concerning observation and "delayed choice" in quantum mechanics. *Physical Review A*, 25(4), 2208–2213.
    - This foundational paper on the quantum eraser experiment introduces the concept of “erasing” which-path information, providing a basis for the experimental methodology in this paper.
6. **Lambert, N., Chen, Y.-N., Cheng, Y.-C., Li, C.-M., Chen, G.-Y., & Nori, F.** (2013). Quantum biology. *Nature Physics*, 9(1), 10–18.
    - An overview of quantum effects in biological systems, discussing phenomena like coherence and superposition in biological processes, supporting the relevance of quantum mechanics in photosynthesis.
7. **van Grondelle, R., & Novoderezhkin, V. I.** (2006). Energy transfer in photosynthesis: Experimental insights and quantitative models. *Nature Physics*, 2(7), 437–447.
    - Reviews experimental techniques and quantitative models for studying energy transfer in photosynthetic systems, with insights into electron transfer pathways and coherence.
8. **Brixner, T., Stenger, J., Vaswani, H. M., Cho, M., Blankenship, R. E., & Fleming, G. R.** (2005). Two-dimensional spectroscopy of electronic couplings in photosynthesis. *Nature*, 434(7033), 625–628.
    - Discusses the use of two-dimensional spectroscopy for studying coherence in photosynthetic systems, providing methodological insights applicable to time-resolved measurements in the proposed experiment.
9. **Fassioli, F., Olaya-Castro, A., & Scholes, G. D.** (2012). Coherent energy transfer under incoherent light conditions. *The Journal of Physical Chemistry Letters*, 3(24), 3136–3142.
    - Examines coherence under natural lighting, which is relevant for understanding how quantum effects might persist in biological systems exposed to noisy environments.

10. **Lloyd, S.** (2011). Quantum coherence in biological systems. *Journal of Physics: Conference Series*, 302, 012037.
    - Discusses the theoretical aspects of quantum coherence in biological systems, reinforcing the framework that quantum effects may contribute to biological functionality.
11. **Brookes, J. C.** (2017). Quantum effects in biology: Golden rule in enzymes, olfaction, photosynthesis and magnetodetection. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 473(2201), 20160822.
    - A review of the evidence for quantum effects across various biological systems, relevant for supporting the plausibility of coherence in photosynthetic electron transfer.
12. **Marcus, R. A., & Sutin, N.** (1985). Electron transfers in chemistry and biology. *Biochimica et Biophysica Acta (BBA) - Reviews on Bioenergetics*, 811(3), 265–322.
    - Foundational paper on the mechanisms of electron transfer in biology, covering factors that influence pathway selectivity and reaction kinetics in RCs.
13. **Chenu, A., & Scholes, G. D.** (2015). Coherence in energy transfer and photosynthesis. *Annual Review of Physical Chemistry*, 66, 69–96.
    - Reviews coherence in energy transfer, providing a comprehensive look at how quantum coherence may be preserved and utilized in photosynthetic systems.
14. **Mancal, T., & Valkunas, L.** (2010). Exciton dynamics in photosynthetic complexes. *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 1797(9), 1507–1516.
    - Discusses the dynamics of exciton transfer in photosynthetic complexes, with relevance to quantum superposition and pathway selection in RCs.

15. **Mohseni, M., Rebentrost, P., Lloyd, S., & Aspuru-Guzik, A.** (2008). Environment-assisted quantum walks in photosynthetic energy transfer. *The Journal of Chemical Physics*, 129(17), 174106.
    - This paper models how environmental interactions can assist coherence in photosynthetic energy transfer, providing insights into coherence maintenance in complex biological environments.
16. **Prokhorenko, V. I., Nagy, A. M., Waschuk, S. A., Brown, L. S., Birge, R. R., & Miller, R. J. D.** (2006). Coherent control of retinal isomerization in bacteriorhodopsin. *Science*, 313(5791), 1257–1261.
    - Demonstrates coherent control in a biological system, supporting the feasibility of manipulating coherence and pathway selection in biological samples.
17. **Wolynes, P. G.** (2009). Quantum coherence and entanglement in biomolecular function. *Proceedings of the National Academy of Sciences*, 106(16), 6649–6650.
    - Reviews theoretical perspectives on coherence and entanglement in biomolecular processes, laying the groundwork for coherence-based experiments in biological systems.

These references provide foundational knowledge on photosynthetic electron transfer, quantum coherence, and experimental approaches to studying quantum effects in biology, supporting the rationale and methods proposed in the experiment.


Quantom wicherise
B
A
B
B
B
A
A
Quantum transefecrs